

Fig. 1

6

Clone 1 cDNA: malate dehydrogenase 2

Clone ID: 1500012M15 3

DNA sequence

..... 4

.....

.....

.....

.....

Clone 2 cDNA: isocitrate dehydrogenase (NAD)

Clone ID: 1500012E04

DNA sequence

.....

.....

.....

.....

.....

Clone 3 cDNA: isocitrate dehydrogenase (NADP)

Clone ID: E030024J03

DNA sequence

.....

.....

.....

.....

.....

Clone 4 cDNA: oxoglutarate dehydrogenase (lipoamine)

Clone ID: E430020N12

DNA sequence

.....

.....

.....

.....

.....

**Procedures**

To amplify cDNA Inserts, cut out the 4 mm X 4 mm area spotted with DNA, place it into a PCR tube, add 25  $\mu$  of water, centrifuge the resulting solution, and then initiate PCR cycle. The spot contains Taq DNA polymerase, an aptamer against the Taq DNA polymerase, trehalose, a PCR buffer, PCR primers (-21M13: 5'-TGTAACGACGCGCCAGT-3', 1233-Rv:5'-AGCGGATAACAATTCACACAGGA-3'), each of dATP, dGTP, dCTP and dTTP, and MgCl<sub>2</sub>. After centrifuging the resulting solution, the PCR cycle is initiated. PCR cycles comprise 2 min at 94°C, 29 cycles of denaturing (94°C, 1 min), annealing (60°C, 1 min) and extension (68°C, 75 sec), and 15 min at 74°C.

Clone 1

Clone 2

Clone 3

Clone 4




Fig. 2





Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8






Fig. 9



<p><b>DNA Reagent</b></p> <p><b>Contents of kit</b></p> <p>-----</p> <p>-----</p> <p><b>Directions for use</b></p> <p>-----</p> <p>-----</p> <p>-----</p> <p>-----</p> <p>-----</p> <p>-----</p> <p>-----</p>	<p><b>Cautions in use and handling</b></p> <p>(1) -----</p> <p>(2) -----</p> <p>(3) -----</p> <p>(4) -----</p> <p><b>Customer inquiry</b></p> <p>K.K. Dnaform</p> <p><b>Customer Support Center TEL:</b></p> <p><b>Hours of operation: Mon – Fri</b></p> <p><b>(Closed on Sat + Sun and national holidays.)</b></p>
---	--



Fig. 10



Fwd T7 loxP Sfi I Sal I \*\*\*\*\* Bam HI Sfi I T3 Rev  
TGTAAAACGACGGCCAGTGAATTGT AATACGACTCACTATAGGGCGAATTGGAGC  
TCCACCGCGGTGCCCCCCCCATAACTTCGTATAGCATAATTATACGAAATTATG  
GATCAGGCCAAATCGGCCGAGCTCGAATTCGTCGAC\*\*\*\*\*  
 \*\*\*\*\*  
GGATCCCGCCATAAGGGCC TGATCCTTCGAGGGGGGGCCCGGTACCAGCTTTTGT  
TGCTTTAGTGAGGGTTAATTTCGAGCTTGGCGTAATCATGTCATAGCTGTTTC  
CTGTGTGAAATTGTT

Fig. 11

	<b>cDNA: malate dehydrogenase</b>
	<b>Clone ID: 1500012M15</b>
	<b>DNA sequence</b>
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	.....
	<b>Amino acid sequence</b>
	.....
	.....
	.....
	.....
	.....
	<b>Primer</b>
	—21M13 : 5'-TGTAACGACGGCCAGT-3'
	1233-Rv : 5'-AGCGGATAACAATTTACACAGGA-3'
	malate dehydrogenase cDNA
	primers

Fig. 12




Fig. 13




Fig. 14

Lane 1: primer set 1, aptamer (-)  
Lane 2: primer set 2, aptamer (-)  
Lane 3: size marker  
Lane 4: primer set 1, aptamer(+)  
Lane 5: primer set 2, aptamer(+)




Fig. 15

Clone 1 cDNA: malate dehydrogenase  
Clone ID: 1500012M15

DNA sequence

.....  
.....  
.....  
.....  
.....

Clone 2 cDNA: isocitrate dehydrogenase (NAD)  
Clone ID: 1500012E04

DNA sequence

.....  
.....  
.....  
.....  
.....

Clone 3 cDNA: isocitrate dehydrogenase (NADP)  
Clone ID: E030024J03

DNA sequence

.....  
.....  
.....  
.....  
.....

Clone 4 cDNA: oxoglutarate dehydrogenase (lipoamine)  
Clone ID: E430020N12

DNA sequence

.....  
.....  
.....  
.....  
.....

Clone 1



Clone 2



Clone 3



Clone 4



Fig. 16




Fig. 17



BEST AVAILABLE COPY